

ATTACHMENT 3

2770834

EXPIRES

3 1284313105 7

0744

148191

PAIN RELIEVER/FEVER REDUCER

# BAYER

GENUINE ASPIRIN

## CHILDREN'S

### 81mg Chewable

ORANGE FLAVOR

36 CHEWABLE TABLETS 81 mg

Bayer Corporation  
P.O. Box 1310  
Morton, NJ 07962-1310 USA

USE ONLY IF SEAL UNDER BOTTLE CAP WITH  
WHITE Bayer Corporation PAIN IS MARCH

**Drug Facts (continued)**

**Other Information**

- Save carton for full directions and warnings.
- Store at room temperature.

**Inactive Ingredients**

Deutose, FD&C Yellow #6 Aluminum Lake, Flavor, Saccharin Sodium, Starch

**Questions or comments?**

1-800-331-4636 or [www.bayeraspirin.com](http://www.bayeraspirin.com)

### Drug Facts

**Active ingredient (in each tablet)** Aspirin 81 mg **Purpose** Pain Reliever/Fever Reducer

**Uses:** For the temporary relief of

- minor aches, pains, and headaches
- to reduce fever associated with colds, sore throats, and teething

#### Warnings

**Reye's syndrome:** Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

**Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding.

**Do not use** • If you are allergic to aspirin • for at least 7 days after tonsillectomy or oral surgery

#### Drug Facts (continued)

**Ask a doctor before use if you have**

- a child experiencing arthritis pain
- stomach problems (such as heartburn, ulcers, or acid pain) that persist or recur

**Ask a doctor or pharmacist before use if you have**

- anticoagulation (thinning the blood) or are taking an anticoagulation drug for arthritis

**Stop use and ask a doctor if**

- pain or fever persists or gets worse
- new or unexpected symptoms occur
- sore throat is severe, persists, or is accompanied by fever, headache, rash, nausea, or vomiting

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems of the unborn child or complications during delivery.**

#### Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- Do not use if pain persists for more than 10 days (for adults) or 5 days (for children).
- Do not use if fever persists for more than 3 days. • Do not exceed recommended dosage.
- To be administered only under adult supervision. • Drink water with each dose.

Age (years)	Weight (lbs)	Dosage	Maximum Dosage
Children under 3	Under 32		Consult a doctor.
3 to under 4	32 to 35	2 tablets	
4 to under 6	36 to 45	3 tablets	
6 to under 9	46 to 65	4 tablets	Repeat every 4 hours while symptoms persist up to a maximum of 5 doses in 24 hours or as directed by a doctor
9 to under 11	66 to 76	4 to 5 tablets	
11 to under 12	77 to 83	4 to 6 tablets	
Adults and children 12 years and over		5 to 8 tablets	